

On a Linear Property of Bilinearly Defined Sets in $\mathbb{R}^n$

LEO LIBERTI¹

*Centre for Process Systems Engineering
Imperial College of Science, Technology and Medicine
London SW7 2BY
United Kingdom*

9 July 2002

Abstract

Certain types of bilinearly defined sets in $\mathbb{R}^n$ exhibit a higher degree of linearity than what is apparent by inspection.

Let $n, m \in \mathbb{N}$ such that $n < m$; $w, x \in \mathbb{R}^n$; $y \in \mathbb{R}$; J be an index set of size $n - m$ such that for all $j \in J$ we have $j \leq n$; $A = (a_{ij})$ an $m \times n$ matrix having rank m ; $b \in \mathbb{R}^m$;

$$C = \{(x, w, y) \mid Ax = b \wedge \forall j \leq n (w_j = x_j y)\}; \quad (1)$$

$$R_J = \{(x, w, y) \mid Ax = b \wedge Aw - by = 0 \wedge \forall j \in J (w_j = x_j y)\}. \quad (2)$$

Note that since $y \in \mathbb{R}$, the product by is simply $(b_1 y, \dots, b_m y)^T$. We shall show that there is at least a J such that $C = R_J$. This in effect means that m of the n bilinear terms that define the set C can be replaced by the linear system $Aw - by = 0$. Thus the set C has a higher degree of linearity than what is apparent by inspection of eqn. (1).

Theorem

$\exists J (C = R_J)$.

Proof. If $(x, w, y) \in C$, then by definition it satisfies $Ax = b$; multiplying this system by y we obtain $yAx - by = 0$, and since $w_j = x_j y$ for all $j \leq n$ we have $Aw - by = 0$. Hence $(x, w, y) \in R_J$ for all J . Now because $\text{rk}(A) = m$, by applying Gaussian elimination we can find a permutation $\pi \in S_n$ (the symmetric group of order n), a vector $b' \in \mathbb{R}^m$ and a matrix $A' = (A'_0 \mid A'_1)$, where:

$$A'_0 = \begin{pmatrix} a'_{11} & a'_{12} & \cdots & a'_{1m} \\ 0 & a'_{22} & \cdots & a'_{2m} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & a'_{mm} \end{pmatrix}, \quad A'_1 = \begin{pmatrix} a'_{1,m+1} & \cdots & a'_{1n} \\ a'_{2,m+1} & \cdots & a'_{2n} \\ \vdots & \ddots & \vdots \\ a'_{m,m+1} & \cdots & a'_{mn} \end{pmatrix},$$

A'_0 being an $m \times m$ square matrix of rank m , such that $A'\pi(x) = b'$. Let $J' = \{m+1, \dots, n\}$, $J = \pi^{-1}(J')$, and $(x, w, y) \in R_J$. Since $b = Ax$ we can substitute it into the system $Aw - by = 0$ to get $Aw - yAx = 0$, i.e. $A(w - xy) = 0$. This is an underdetermined linear system with m equations and n variables

$$\forall i \leq m \quad \left(\sum_{j=1}^n a_{ij}(w_j - x_j y) = 0 \right),$$

¹E-mail: l.liberti@ic.ac.uk.

which (by Gaussian elimination) has the same solution space as the system $A'\pi(w - xy) = 0$. Since for all $j \in J$ we have, by definition of R_J , $w_j = x_j y$, this is in turn equivalent to

$$A'_0 \bar{\pi}(w - xy) = 0 \quad (3)$$

where $\bar{\pi}$ is a restriction of π to $\{1, \dots, n\} \setminus J$ (so that $\bar{\pi}(j) = \pi(j)$ only if $j \notin J$). The system (3) is square with full rank m and thus has a unique solution $\bar{\pi}(w - xy) = 0$, i.e.

$$\forall j \notin J \ (w_j = x_j y).$$

Hence (x, w, y) satisfies the definition of C , i.e. $R_J \subseteq C$. □

The geometrical implications of this theorem are that the intersection in $\mathbb{R}^n$ of a set of bilinear terms like those described above is a hypersurface containing a degree of linearity. By exploiting this linearity we are able to replace some of the bilinear terms with linear terms. This fact is useful in a number of applications, e.g. linearly constrained bilinear programming.